

SENTRY

Dual Action Flea & Tick Collar for Dogs

READ ENTIRE LABEL BEFORE EACH USE • USE ONLY ON DOGS
PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION

HAZARDS TO HUMANS: Harmful if swallowed. Harmful if absorbed through the skin. Causes moderate eye irritation. Avoid contact with skin, eyes or clothing. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove and wash contaminated clothing before reuse. **HAZARDS TO DOMESTIC ANIMALS:** Collar is for external use only, not to be taken internally. Some animals may become irritated by any collar. If this occurs, remove collar. If condition persists, consult a veterinarian. Consult a veterinarian before using this product on debilitated, aged, medicated, pregnant or nursing animals. Do not use on pets under 12 weeks of age. Sensitivities may occur after using ANY pesticide product for pets. If signs of sensitivity occur remove collar and bathe your pet with mild soap and rinse with large amounts of water. If signs continue, consult a veterinarian immediately.

Notice: This product contains a chemical (o-Isopropoxyphenyl methylcarbamate) known to the State of California to cause cancer.

FIRST AID

IF SWALLOWED:	• Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by the poison control center or doctor. • Do not give anything by mouth to an unconscious person.
IF ON SKIN OR CLOTHING:	• Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
IF IN EYES:	• Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.

HOT LINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-781-4738 for emergency medical treatment information.

NOTE TO PHYSICIAN/VETERINARIAN:

Dust released by this collar is a cholinesterase inhibitor. Atropine is antidotal. Do not use this product on animals simultaneously or within 30 days or after treatment with or exposure to cholinesterase inhibiting drugs, pesticides or chemicals. However, flea and tick collars may be immediately replaced.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

- Do not open until ready to use. Dust will form on collar during storage. Do not get dust in mouth or eyes. Take the collar out of the packaging and stretch to full length. The tabs that held the collar in place will fall off when stretching; pull on any remaining tabs to remove them.
- Bend the buckle in half between the holes in the buckle, so that you can easily pass the end of the collar through both holes.
- Place the collar around your pet's neck. Feed the collar through both holes, notches facing out. Fit to the neck of the dog. The collar must be worn loosely to allow for growth of the pet and to permit the collar to move about the neck. Generally, a properly fitted collar is one that, when fastened, will snugly slide over pet's head, and two fingers can be placed between the collar and pet's neck. Leave 1 or 2 extra inches for collar adjustment, cut off remaining excess collar and dispose of in the trash.
- Wash hands after installing collar on pet. Replace collar every 6 months, or sooner if needed.

STORAGE AND DISPOSAL

PESTICIDE STORAGE: Store only in original container in a cool, dry area inaccessible to children and pets. Protect from freezing.
PESTICIDE DISPOSAL AND CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container. If empty: Place in trash or offer for recycling, if available. If partly filled: Call your local solid waste agency for disposal instructions. Dispose of expired collar in trash.

In Case of Emergency, call: 1-800-781-4738
Non-Emergency, call: 1-800-224-PETS (7387)

EPA Reg. No. 2517-61
EPA Est. No. 42836-TX-1^a
EPA Est. No. 2517-NE-Q2^a

Superscript is the first letter in the lot code

If you have questions or comments about this product, please write:
SENTRY Consumer Response, P.O. Box 540399, Omaha, NE 68154-0399
Distributed by: Sergeant's Pet Care Products, Inc.
Omaha, NE 68154-0399

P004426 ANRC

SENTRY

Dual Action Flea & Tick Collar for Dogs

Fits necks to
23"

Fits necks to
23"



Fits necks to
23"

Fits necks to
23"

Dual Action Flea & Tick Collar for Dogs
SENTRY

SENTRY

Dual Action Flea & Tick Collar for Dogs

- Kills Fleas and Ticks for up to 6 months
- Kills the Ticks that may carry Lyme Disease



RIGHT FIT design

DO NOT ALLOW CHILDREN TO PLAY WITH COLLAR

CAUTION SEE BACK LABEL PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

NET WEIGHT: .97 oz (27.5 g)

ACTIVE INGREDIENTS:

Propoxur	9.00%
d-(cis,trans) phenothrin*	3.00%
N-Octyl bicycloheptene dicarboximide	3.00%
OTHER INGREDIENTS:	85.00%
TOTAL	100.00%

*cis/trans isomer ratio: Max. 25% (+/-) cis and Min. 75% (+/-) trans

Dual Action Flea & Tick Collar for Dogs
SENTRY

UNCOATED AREA

EPA Reg. No. 2517-61



0 73091 01992 3