10324-155-5741 HDQ Neutral 20211008 45 5741 .pdf

HDQ Neutral®

This product has been designed specifically where housekeeping is of prime moortance. This product when used as directed is formulated for use on nard, non-porous surfaces such as sealed floors, walls, countertops, tables, chairs, bed frames and bed railings.

DISINFECTION PERFORMANCE: This product kills the following bacteria in minutes at 1 oz. per gal. of 400 ppm hard water (660 ppm active) and 6 soil on hard, non-porous surfaces

inetobacter baumannii, Acinetobacter Iwoffi, Bordetella bronchiseptica, Citrobacter freundii, Enterobacter agglomerans, Enterobacter cloacae, scherichia coli, Escherichia coli 0157:H7, Escherichia coli (Extended pectrum B-Lactamase) (ESBL), Escherichia coli (Tetracycline Resistant), terococcus faecalis. Enterococcus hirae. Fusobacterium necrophorum. (lebsiella aerogenes, Klebsiella oxytoca, Klebsiella pneumoniae, Klebsiella pneumoniae (Carbapenem-Resistant). Listeria monocytogenes Micrococcus luteus, Pasturella multocida, Proteus vulgaris, Pseudomonas eruginosa, Pseudomonas aeruginosa (Tetracycline Resistant), idomonas cepacia, Salmonella entèrica, Salmonella enterica serotype ıllorum, Salmonella typhi, Salmonella typhimurium, Serratia marcescens, higella flexneri, Shigella sonnei, Staphylococcus aureus, Staphylococcus ireus (Community Associated Methicillin Resistant) (CA-MRSA). aphylococcus aureus (Methicillin Resistant) (MRSA), Staphylococcus dermidis (Ampicillin, Cefazolin, Oxacillin Resistant), Staphylococcus idermidis, Streptococcus agalactiae, Staphylococcus haemolyticus, reptococcus pneumoniae (Penicillin Resistant). Streptococcus pyogenes. reptococcus mutans, Enterococcus faecalis (Vancomycin Resistant) RE), Staphylococcus aureus (Vancomycin Intermediate Resistant) (VISA)

VIRUCIDAL* PERFORMANCE: This product kills the following viruses in) minutes at 1 oz. per gal. of 400 ppm hard water (660 ppm active) and 6 soil on hard, non-porous surfaces: Avian Influenza A (H3N2) Virus, Avian Influenza A (H5N1) Virus, Cytomegalovirus, Coronavirus (SARS) associated), Hepatitis B Virus, Hepatitis C Virus, Herpes Simplex Virus Type Herpes Simplex Virus Type 2, Human Coronavirus, Influenza A Virus 3N2), Human Immunodeficiency Virus Type 1 (HIV 1), Influenza A Virus H1N1), Respiratory Syncytial Virus, SARS-CoV-2, Vaccinia Virus.

Vibrio cholera, Yersinia enterocolitica

ANIMAL PREMISE VIRUCIDAL* PERFORMANCE: This product kills the ollowing viruses in 10 minutes at 1 oz. per gal. of 400 ppm hard water (660 pm active) and 5% soil on hard, non-porous surfaces: Avian Infectious onchitis Virus, Avian Influenza A Virus, Canine Coronavirus, Canine temper Virus, Feline Picornavirus, Infectious Bovine Rhinotracheitis Virus, Pseudorabies Virus, Swine Influenza A Virus, Transmissible Gastroenteritis Virus

This product kills the following viruses in 10 minutes at 4.5 oz. per gal. of 400 ppm hard water (2970 ppm active) and 5% soil on hard, non-porous surfaces: Canine Parvovirus** (Type 2b), Rabies Virus**

** Indicates that a dilution of 4.5 oz. per gal. of water is required for this

FUNGICIDAL PERFORMANCE: This product kills the following fungi in 10 minutes at 1 oz. per gal. of 400 ppm hard water (660 ppm active) and 5% soil on hard, non-porous surfaces: Candida albicans, Trichopyhton interdigitale (Athlete's foot fungus) (Ringworm fungus)

MILDEWSTATIC PERFORMANCE: This product controls the following mold at 1 oz. per gal. of 400 ppm hard water (660 ppm active) and 5% soil on hard, non-porous surfaces for up to 7 days: Aspergillus niger

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling

Rinse hard, non-porous food contact surfaces with potable water after application of product.

This product is not for use on medical device surfaces

	DILUTION TABLE:	Dilution	Contact Ti
	For Hospital or Medical Environment Disinfection Claims	1 oz. /gal. water	10 minutes
	For General or Broad Spectrum Disinfectant Claims	· ·	10 minutes
,	For Public Health Virucidal* claims	1 oz. /gal. water	10 minutes
	For Animal Virucidal** Claims For Fungicidal Claims	4.5 oz. /gal. water 1 oz. /gal. water	10 minutes 10 minutes

For Mold and Mildew Claims 1 oz. /gal. water Reapply every 7 days FOR USE AS A ONE-STEP, GENERAL, HOSPITAL DISINFECTANT, FUNGICIDE VIRUCIDE* DEODORIZER CLEANER

. Pre-clean visibly soiled areas.

Sold by: SPARTAN CHEMICAL COMPANY, INC., 1110 Spartan Drive, Maumee, OH 43537 USA / 1-800-537-8990 /For medical emergencies, call 888-314-6171 / www.spartanchemical.com

- 2. Apply use solution of 1 oz. of this product per gal. of water (or equivalent use dilution) to disinfect hard, non-norous, non-food contact surfaces. with a sponge, brush, cloth, mop, by immersion, auto scrubber. mechanical spray device, hand pump trigger spray device, coarse trigger spray device. For spray applications, spray 6-8 inches from surface. Do not breathe spray.
- Treated surfaces must remain visibly wet for 10 minutes
- 4. Allow to air dry. Rinsing of floors is not necessary unless they are to be
- 5. Prepare a fresh solution daily or when visibly dirty.

*KILLS HIV HRV AND HCV ON PRE-CLEANED HARD NON-POROUS SURFACES/OBJECTS PREVIOUSLY SOILED WITH BLOOD/BODY FLUIDS in health care settings or other settings in which there is an expected

likelihood of soiling of hard, non-porous surfaces/objects with blood or hody fluids and in which the surfaces/objects likely to be soiled with blood or body fluids can be associated with the potential for transmission of Human Immunodeficiency Virus Type 1 (HIV-1) (associated with AIDS), Hepatitis B Virus (HBV) and Hepatitis C Virus (HCV).

SPECIAL INSTRUCTIONS FOR CLEANING AND DECONTAMINATION AGAINST HIV-1. HBV and HCV ON SURFACES/OBJECTS SOILED WITH BLOOD/BODY FLUIDS

Personal Protection: Wear protective latex gloves, gowns, masks and eve

Cleaning Procedure: Blood and other body fluids must be thoroughly cleaned from hard, non-porous surfaces and objects before application of

Disposal of Infectious Materials: Blood and other body fluids, cleaning materials and clothing must be autoclaved and disposed of according to federal, state and local regulations for infectious waste disposal. Contact Time: Allow hard, non-porous surfaces to remain visibly wet for 10

TO CONTROL MOLD/MILDEW: Pre-clean visibly soiled areas. Prepare use solution by adding 1 oz. of this product per gal. of water (or equivalent use dilution). Apply use solution to hard, non-porous surfaces which will effectively inhibit the growth of mold and mildew and their odors. Repeat treatment every seven days, or more often if new growth appears.

ANIMAL PREMISES: Prior to use of this product, remove all animals and feeds from premises, animal transportation vehicles, and enclosures. Remove all litter, droppings and manure from floors, walls and surfaces of

barns, pens, stalls, chutes and other surfaces of facilities and fixtures occupied or traversed by animals. Empty all troughs, racks and other feeding and watering appliances. Thoroughly clean surfaces with soap or detergent and rinse with water.

FOR USE AS AN ANIMAL PREMISE DISINFECTANT/VIRUCIDE*:

Prepare a use solution of 1 oz. of this product per gal. of water (660 ppm active) for effectiveness against designated animal viruses on the

FOR USE AS AN ANIMAL PREMISE DISINFECTANT/VIRUCIDE**:

Prepare a use solution of 4.5 oz. of this product per gal. of water (2970 ppm active) for effectiveness aganst designated animal viruses on the

Apply use solution to disinfect hard, non-porous surfaces with a sponge brush cloth by immersion mechanical spray device hand pump trigger spray device, coarse trigger spray device. For spray applications, spray 6-8 inches from surface. Do not breathe spray. Immerse all halters and other types of equipment used in handling and restraining animals, as well as forks, shovels, and scrapers used for removing litter and manure in the use solution. Treated surfaces must remain visibly wet for 10 minutes. Ventilate buildings, coops and other closed spaces. Do not house animals or employ equipment until treatment has been absorbed, set or dried. Thoroughly scrub all treated feed racks, troughs, automatic feeders, fountains and waterers and other treated equipment which can contact food or water with soap or detergent, and rinse with potable water before reuse.

GENERAL DEODORIZATION: To deodorize, apply 1 - 2 oz. of this product per gal. of water to hard, non-porous surfaces. Rinse or allow to air dry. Rinse with potable water after use on surfaces that come in

STORAGE AND DISPOSAL

o not contaminate water, food, or feed by storage and disposal.

PESTICIDE STORAGE: Store only in original container. Keep this product under locked storage sufficient to make it inaccessible to children or persons

PESTICIDE DISPOSAL: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture or rinsate is a violation of Federal w. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the rdous Waste Representative at the nearest EPA Regional Office for guidance.

ONTAINER HANDLING: Non-Refillable Container. Do not reuse or refill this container, Triple rinse container (or equivalent) promptly after emptying. iple rinse as follows: Fill the container 1/4 full with water and recap. Shake for 10 seconds. Drain for 10 seconds after the flow begins to drip. Follow esticide Disposal instructions for rinsate disposal. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration.

K-20 SCC 4/21



ACTIVE INGREDIENTS:

Ammonium Chloride.....

Alkvl (50% C14, 40% C12,

10% C16) Dimethyl Benzyl

Ammonium Chloride.....

EPA Reg. No. 10324-155-5741

SI USTED NO ENTIENDE LA

PARA QUE SE LA EXPLIQUE A

USTED EN DETALLE

IT TO YOU IN DETAIL.

ETIQUETA, BUSQUE A ALGUIEN

IF YOU DO NOT UNDERSTAND THE

LABEL, FIND SOMEONE TO EXPLAIN

EPA Est. No. 5741-0H-1

..100.009

Didecyl Dimethyl

INGREDIENTS:.

Cleans. Disinfects and Deodorizes

Disinfectant, Mildewstat, Fungicide, Virucide* Effective in hard water up to 400 ppm hardness in the presence of 5% soil. Effective against 2019 SARS-CoV-2 (COVID-19 virus)

KEEP OUT OF REACH OF CHILDREN DANGER PELIGRO

FIRST AID

In case of emergency, call a poison control center or doctor for treatment advice. Have the product container or label with you when calling a poison control center or doctor, or going for treatment

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes,

IF ON SKIN: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes.

IF SWALLOWED: Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person. Call a poison control center or doctor immediately for treatment advice. IF INHALED: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-

NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

DANGER. Corrosive. Causes irreversible eye damage and skin burns. Harmful if swallowed or absorbed through the skin. Do not get in eyes, on skin, or on clothing. Wear goggles or face shield and chemical resistant gloves and protective clothing when handling. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove contaminated clothing and wash

This product is toxic to fish, aquatic invertebrates, oysters and shrimp.

#1202 NET CONTENTS: 1 U.S. Gallon (3.79 Liters)